

Veterinär-Anatomisches Institut
der Vetsuisse-Fakultät Universität Zürich

Direktor ad interim: Prof. Dr. B. Spiess

Arbeit unter Leitung von Prof. Dr. P. Wild

**The Significance of US3 Kinase in Herpes Simplex Virus 1
Envelopment as Revealed by High Resolution Electron Microscopy**

Inaugural-Dissertation

zur Erlangung der Doktorwürde der
Vetsuisse-Fakultät Universität Zürich

vorgelegt von

Sabine Leisinger
Tierärztin
von Davos Dorf, GR

genehmigt auf Antrag von

Prof. Dr. P. Wild, Referent

Prof. Dr. M. Ackermann, Korreferent

Zürich 2006

CONTENT

1. Abstract	1
2. Introduction	2
2.1. Herpes Simplex Virus Type 1	2
2.2. The Theories of Viral Egress	2
2.2.1. Theory of Envelopment, De-Envelopment and Re-Envelopment	2
2.1.2. Theory of Egress by Vesicle Formation	3
2.1.3. Theory of Two Diverse Pathways	4
2.3. Significance of Tegument and Envelope Proteins	6
2.4. Significance of US3	8
2.5. Aim of the Study	10
3. Material and Methods	12
3.1. Cells and Viruses	12
3.2. Infection of Cells	12
3.3. Brefeldin A	12
3.4. Low-Temperature Transmission Electron Microscopy	13
3.5. Morphometric Analysis	13
3.6. Immunolabeling	14

4. Results	15
4.1. Nucleus	15
4.1.1. Nuclear Membranes	15
4.1.2. Nuclear Capsids	24
4.1.3 Nuclear Pores	26
4.2. Cytoplasm	28
4.2.1. RER-Golgi Complex	28
4.2.2. Distribution of Phenotypes	30
4.3. Virus Yields	33
4.4. Effects of Brefeldin A on HSV-1 ΔUS3 Envelopment	33
5. Discussion	40
6. References	46
7. Acknowledgements	52
8. Curriculum Vitae	53

1. Abstract

Capsids of herpesviruses acquire tegument and envelope by budding at the inner nuclear membrane. US3, a protein kinase, plays an important role in envelopment together with a complex of viral proteins UL31 and UL34. To clarify the significance of US3 Vero cells were infected with a US3 deletion mutant of herpes simplex virus 1. High resolution electron microscopy revealed dramatic folding of the inner nuclear membrane in combination with deposition of a dense substance at the luminal layer, accumulation of capsids within the nucleus and the nuclear periphery, accumulation of virions containing a dense envelope within the perinuclear space, dilation of nuclear pores and a few capsids within the cytoplasm. These alterations compared to wildtype infected cells indicate retardation of budding and of release of virions out of the perinuclear space, and perturbation of the nuclear envelope. Exposure to Brefeldin A, which disassembles the Golgi complex, led to a reduction of membrane foldings, and a shift of virions from the perinuclear space into adjacent RER cisternae. US3 is hence suggested to be involved in regulation of distribution of membrane constituents needed for budding and intraluminal virus transportation acting possibly at the level of the Golgi complex. Yields of infectious virus was equal to that of wild type virus indicating that capsids acquire the requirements for infectivity, i.e. tegument and glycoproteins, in the course of budding at the inner nuclear membrane.

2. Introduction

2.1. Herpes Simplex Virus Type 1 (HSV-1)

Herpesviruses are composed of four morphological distinct substructures: the inner core containing the viral genomic DNA, the icosahedral capsid built of 162 capsomers, the tegument surrounding the capsid, and the viral envelope with embedded glycoproteins. The way of assembly of these different structures is highly complex. Capsids formed within nuclei are transported to the nuclear periphery. Translocation through the nucleocytoplasmic barrier, envelopment, tegumentation and intracellular transportation are controversially discussed for decades that has led to three main theories.

2.2. The Theories of Viral Egress

2.2.1. Theory of Envelopment, De-Envelopment and Re-Envelopment

This theory assumes that capsids are enveloped by budding at the inner nuclear membrane, de-enveloped by fusion of the primary envelope with the outer nuclear membrane, and finally, re-envelopment at cytoplasmic membranes (Stackpole 1969). The first budding step in herpesvirus maturation at the inner nuclear membrane provides the capsid with a primary envelope (Darlington and Moss 1968; Stackpole 1969). The net result of the budding event is the formation of primary enveloped virions located in the perinuclear space. Capsids of primary virions are translocated into the cytoplasm after fusion of the primary envelope with the outer nuclear membrane (Stackpole 1969). Primary virions as observed in cells infected with the alphaherpesviruses HSV-1, pseudorabies virus (PrV), equine herpes virus 1 (EHV-1) and infectious laryngotracheitis

virus differs in morphology from mature extracellular virions in that the spikes, which are prominent on mature virions, are not detectable by electron microscopy in virions within the perinuclear space (Granzow 2001). The appearance of the primary tegument differs from that in mature virions in that it is closely apposed to the primary envelope in a characteristic electron dense sharply bordered ring structure (Granzow 2001). Primary and secondary virions differ not only in morphology but also biochemically. Differences in the phospholipid composition between mature HSV-1 virions and the nuclear membrane indicated that the latter may not form the final envelope (van Genderen, Brandimarti et al. 1994). The UL31 and UL34 gene products have been shown to be constituents of primary enveloped virions but not of mature virus particles (Klupp, Granzow et al. 2000; Fuchs, Klupp et al. 2002; Reynolds, Wills et al. 2002). Other tegument and envelope proteins present in mature virions have not been detected in primary virus particles (Granzow, Klupp et al. 2004; Mettenleiter 2004). Budding of herpesvirus capsids in the cytoplasm after translocation through the nuclear membrane has been observed early in electron microscopic studies (Siminoff and Menefee 1966; Stackpole 1969; Roffman, Albert et al. 1990). Recently, it has been propagated that this sequential envelopment–de-envelopment–re-envelopment theory (Skepper, Whiteley et al. 2001) constitutes the prototypic herpesvirus maturation pathway (Mettenleiter 2002). Tegumentation may start at two different sites. One site is the translocated capsid into the cytoplasm, the other is the future envelopment site (Mettenleiter 2002), the trans Golgi network.

2.2.2. Theory of Egress by Vesicle Formation

This theory assumes that perinuclear virions exit the perinuclear space by vesicle formation at the outer nuclear membrane (Campadelli-Fiume, Farabegoli

et al. 1991). Virions are then transported within these vesicles to the Golgi apparatus for further maturation (Morgan, Ellison et al. 1954; Poliquin, Levine et al. 1985; Lee, Bao et al. 1987; Torrisi, Di Lazzaro et al. 1992; Gershon, Sherman et al. 1994; Ward, Campadelli-Fiume et al. 1994; Di Lazzaro, Campadelli-Fiume et al. 1995) and then to the plasma membrane where virions are released into the extra cellular space by exocytosis.

2.2.3. Theory of Two Diverse Pathways

Recently, the uneconomic de-re-envelopment theory has been challenged on the basis of the following observations (Wild, Schraner et al. 2002; Leuzinger, Ziegler et al. 2005; Wild, Engels et al. 2005):

i) Virions are located within RER cisternae indicating transportation of virions from the perinuclear space into RER cisternae, ii) the presence of virions within Golgi cisternae, iii) connectivity between RER and Golgi cisternae suggesting intraluminal transportation of virions from the RER into Golgi cisternae, iv) indications for packaging at lateral Golgi cisternae that results in vacuoles containing one or several virions, v) indications for budding at the outer nuclear membrane, membranes of RER, Golgi complex and Golgi derived vacuoles, vi) budding of capsids at narrow Golgi cisternae resulting in small concentric vacuoles containing a single virion, vii) impaired nuclear pores through which capsids may gain direct access to the cytoplasm. On the basis of these facts a novel theory was developed that includes two diverse routes.

Nuclear envelopment:

Capsids bud through the inner nuclear membrane, and are then transported from the perinuclear space via RER cisternae into Golgi cisternae for packaging into transport vacuoles of various size containing one or more virions (Wild,

Schraner et al. 2002; Leuzinger, Ziegler et al. 2005; Wild, Engels et al. 2005). Fusion of the viral envelope with cell membranes of the compartments the virion is transported through is considered likely to be prevented by proteins of an unknown nature that are condensed at the viral envelope. In addition, these substance or substances facilitate transportation. These “antifusion proteins and/or transport proteins” are speculated to be removed from the envelope in Golgi cisternae but remain in the transport vacuoles at low concentrations to prevent fusion of viral envelopes with the vacuolar membranes.

Cytoplasmic envelopment:

Capsids escape the nucleus via impaired nuclear pores (Leuzinger, Ziegler et al. 2005; Wild, Engels et al. 2005), approach Golgi membranes from the cytoplasmic side and induce budding. Since entire Golgi cisternae are involved in budding, a sphere-like structure comprising two membranes arises. The inner membrane becomes the viral envelope, and the outer one becomes the vacuolar membrane. Fusion of the envelope with the vacuolar membrane is considered likely to be prevented by “antifusion” proteins at high concentrations. Alternatively, capsids escaping the nucleus via impaired nuclear pores can bud at the outer nuclear membrane, RER membranes, membranes of dilated Golgi cisternae, and membranes of Golgi derived vacuoles already containing virions. Virions originating by budding at the outer nuclear membrane and RER membranes need to be transported to Golgi cisternae for packaging.

The fact that capsids acquire both tegument (Roffman, Albert et al. 1990; van Drunen Littel-van den Hurk, Garzon et al. 1995) and an envelope containing glycoproteins (Torrissi, Di Lazzaro et al. 1992; Gilbert and Ghosh 1993; Gilbert, Ghosh et al. 1994) do not support the theory that capsids are de-enveloped by fusion of the primary envelope with the outer nuclear membrane for re-envelopment by Golgi membranes.

2.3. Significance of Tegument and Envelope Proteins

Alphaherpesvirus capsids have to collect more than 15 tegument proteins prior to or concomitantly to acquisition of a lipid envelope containing more than 10 glycoproteins (Steven 1997). Betaherpesvirus need to acquire even more different proteins (Gibson 1996). This assembly is a complex process that is only recently starting to unfold. Only a few of the components of the mature tegument and envelope are conserved which makes it difficult to propose a general pathway of virion formation for all herpesviruses.

For envelopment at the nuclear membrane, two virally encoded proteins have been shown to be crucial. These are the products of the UL31 and UL34 genes of HSV-1 (Chang, Van Sant et al. 1997; Roller, Zhou et al. 2000; Reynolds, Ryckman et al. 2001; Reynolds, Wills et al. 2002). Capsids of primary virions may be translocated into the cytoplasm after fusion of the primary envelope with the outer nuclear membrane (Stackpole 1969). The molecular mechanism of this fusion process is unknown. It is also unclear whether the UL34 protein, which so far is the only membrane protein unequivocally identified as a component of primary virions, is capable of mediating this fusion event either alone or in combination with other viral proteins, e.g. UL31 (Roller, Zhou et al. 2000; Reynolds, Ryckman et al. 2001; Fuchs, Klupp et al. 2002; Reynolds, Wills et al. 2002). These two proteins could also function as transport proteins facilitating transportation of virions into RER cisternae (Leuzinger, Ziegler et al. 2005). UL11 (Baines and Roizman 1992), UL20 (Baines, Ward et al. 1991), UL48 (Mossman, Sherburne et al. 2000), and gK (Foster and Kousoulas 1999; Foster, Rybachuk et al. 2003) have been described to be involved in nuclear egress of HSV-1. It has been proposed that the inner shell of the HSV-1 tegument exhibits icosahedral symmetry and part of it anchored specifically at the vertices (five-fold axis) of the capsid (Zhou, Chen et al. 1999). This inner shell was suggested

to contain the UL36 protein, the largest gene product found in the herpesvirus genome (Baer, Bankier et al. 1984; Chee, Bankier et al. 1990). If the UL36 protein forms the innermost layer of tegument, it has to interact with other tegument proteins for virus maturation to proceed. In agreement with their conservation, the UL36 and UL37 play crucial roles in the tegumentation process. The UL31 and UL34 gene products of PrV are present in virus particles in the perinuclear space but are absent from virions at later stages of assembly (Fuchs, Klupp et al. 2002). In contrast, the UL36, UL37, UL46, UL47, UL48 and UL49 tegument proteins are found only in cytoplasmic enveloped virions and mature PrV particles and are suggested to be presumably not recruited during primary envelopment (Fuchs, Klupp et al. 2002; Klupp, Fuchs et al. 2002; Kopp, Klupp et al. 2002; Granzow, Klupp et al. 2004).

Secondary envelopment has been often observed in close proximity to the Golgi apparatus. Whereas it was proposed that the final envelope is derived from early endosomes (Tooze, Hollinshead et al. 1993) or the endoplasmic reticulum-Golgi-intermediate compartment (Sanchez, Sztul et al. 2000) biochemical and morphological evidence indicate that the vesicles, into which budding occurs, are derived from the Golgi apparatus and most likely belong to the trans-Golgi network (Jones and Grose 1988; Severi, Landini et al. 1988; Whealy, Card et al. 1991; Gershon, Sherman et al. 1994; Zhu, Gershon et al. 1995; Whiteley, Bruun et al. 1999; Granzow 2001; McMillan and Johnson 2001). How these vesicles can increase in size to form finally both the viral envelope and the vacuolar membrane remains mysterious (Leuzinger, Ziegler et al. 2005). It is also unclear how tegumented capsids are directed to the envelopment site and how viral glycoproteins are assembled there. However, recently a conserved virion tegument component, the UL11 protein (MacLean, Clark et al. 1989; Harper and Kangro 1990; Sanchez, Sztul et al. 2000; Loomis, Bowzard et al. 2001; Kopp, Granzow et al. 2003; Schimmer and Neubauer 2003), and a conserved viral

glycoprotein, gM (Klupp, Nixdorf et al. 2000), have been implicated in these (Kopp, Granzow et al. 2004) processes. The UL11 might influence secondary envelopment by directing tegument components (Baines and Roizman 1992; Kopp, Granzow et al. 2003; Silva, Yu et al. 2003) and, conceivably, associated capsids to the budding site whereas gM might act by accumulating envelope glycoproteins at these locations.

Envelopment finally results in an enveloped virion within a secretory vesicle. The vesicle is transported to the plasma membrane where the vesicular membrane fuses with plasma membrane releasing the mature virions into the extracellular space. Little is known about the viral components involved in vesicle transportation and exocytosis. UL20 and gK of HSV-1 and PrV have been implicated in this process (Baines, Ward et al. 1991; Avitabile, Ward et al. 1994; Fuchs, Klupp et al. 1997; Foster and Kousoulas 1999).

2.4. Significance of US3

US3, a protein kinase present in HSV-1 infected cells (Purves, Deana et al. 1986; Purves, Katan et al. 1986), is a product of a viral gene. The enzyme is encoded by the open reading frame designated US3 that maps in the unique sequences of the S component of HSV-1 DNA (Purves, Longnecker et al. 1987; Reynolds, Ryckman et al. 2001). UL34 protein is a substrate for the HSV-1 US3-encoded kinase (Purves, Katan et al. 1986; Frame, Purves et al. 1987; Purves, Spector et al. 1991; Purves, Spector et al. 1992; Roller, Zhou et al. 2000; Shiba, Daikoku et al. 2000; Ye and Roizman 2000; Reynolds, Wills et al. 2002). US3 is required for even distribution of the UL31 and UL34 proteins around the nuclear rim of wild-type infected cells. The HSV-2 US3 kinase has been proposed to phosphorylate the UL34 protein, thereby modulating UL34 function (Purves and

Roizman 1992). Mutants lacking the US3 protein kinase was reported to block transportation of capsids from the nucleus to the perinuclear space (Reynolds, Wills et al. 2002) suggesting that the effect of US3 may be on the integrity of the nuclear envelope and not specifically on the mechanism of capsid transportation. In addition, the US3- encoded kinase has been proposed to play a role in protecting HSV-1 infected cells from virus-induced apoptosis (Leopardi, Van Sant et al. 1997; Galvan and Roizman 1998; Asano, Honda et al. 1999; Jerome, Fox et al. 1999; Munger, Chee et al. 2001).

Interestingly, the PrV UL34 protein is phosphorylated to a similar extent in the presence or absence of US3 protein indicating that the observed phenotype of US3 deletion mutants is not due to differences in UL34 phosphorylation (Klupp, Granzow et al. 2000). In the absence of the PrV US3 protein homologue, large numbers of enveloped virions appear to accumulate within invaginations of the nuclear membrane (Wagenaar, Pol et al. 1995; Klupp 2001). The UL34 gene encodes a type II C-terminally anchored membrane protein (Purves, Spector et al. 1992; Klupp, Granzow et al. 2000) that is located in pseudorabies virus PrV infected cells in both leaflets of the nuclear membrane (Klupp, Granzow et al. 2000).

The UL34 gene of HSV-1 (and PRV) encodes a phosphoprotein that is primarily localized at the nuclear envelope of infected cells. It is a necessary component of an envelopment complex that also includes the UL31 protein (Roller, Zhou et al. 2000; Reynolds, Ryckman et al. 2001; Reynolds, Wills et al. 2002). Localization data and sequence analysis suggest that the UL34 protein is anchored within the inner nuclear membrane by a C-terminal hydrophobic domain leaving the bulk of the protein exposed to the interior of the nucleus (Klupp, Granzow et al. 2000; Reynolds, Ryckman et al. 2001; Reynolds, Wills et al. 2002). There are at least three nonmutually exclusive ways by which UL34 protein could facilitate

nuclear egress (Wagenaar, Pol et al. 1995; Klupp 2001; Reynolds, Wills et al. 2002). First, UL34 could directly mediate envelopment through bridging interactions between the capsid and the inner nuclear membrane. Second, the UL34 protein might direct the localization of other important nuclear egress factors. Third, the nuclear lamina, which lines the interior face of the nuclear envelope, may represent a significant physical barrier to nuclear egress. Herpesvirus infection alters this structure (Scott and O'Hare 2001). UL34 may, therefore, affect the architecture of the nuclear lamina to allow capsids to access the envelopment machinery. Data supporting this function have been reported for the UL34 homologue of murine cytomegalovirus (Muranyi, Haas et al. 2002).

The UL31 gene product is a nuclear matrix associated nucleotidylated phosphoprotein, which in association with the UL34 gene product, localizes to the nuclear rim of HSV-1 infected cells (Chang and Roizman 1993; Blaho, Mitchell et al. 1994; Reynolds, Ryckman et al. 2001; Yamauchi, Shiba et al. 2001; Fuchs, Klupp et al. 2002). The UL31 protein requires UL34 for nuclear membrane targeting (Reynolds, Ryckman et al. 2001; Fuchs, Klupp et al. 2002) whereas the UL34 gene product appears to possess an intrinsic property for targeting to the nuclear membrane (Chang and Roizman 1993). The UL34 protein may thus represent a primary envelope protein whereas the UL31 protein may constitute a primary tegument protein (Mettenleiter 2002).

2.5. Aim of the study

The mutant R7041 was shown to lack the virus induced protein kinase. It was establish that the lack of activity is due the deletion in US3 and not to a fortuitous mutation introduced into the domain of the *tk* gene during the rescue

of the *tk* gene. Concurrently, a mutation in the genes, which partially overlap the *tk* gene, the US3 open reading frame was repaired of recombinant virus R7035 (Purves, Longnecker et al. 1987). R7041 is identical to R7040 except that the 700-base-pair deletion in the *tk* gene was restored by cotransfection of intact R7040 DNA and *Bam*HI-Q DNA cloned as pRB 103 into rabbit skin cells (Purves, Longnecker et al. 1987).

The goal of this study is to clarify the significance of US3 in HSV-1 envelopment. Therefore, cells were infected with HSV-1 Δ US3 for analysis by electron microscopy at improved temporal and spatial resolution, which was achieved by rapid freezing of cells after incubation at 37°C for 17 h to 21 h followed by freeze substitution and embedding. This procedure enabled examinations of virus cell interactions in an *in situ* situation with a delay of less than 15 seconds between removal of the sapphire disks from the incubation chamber and freezing (Wild, Schraner et al. 2002). All processes of envelopment at the nuclear envelope, RER, and Golgi complex could be readily visualized due to the well preserved ultrastructure. Close examination of serial sections revealed a dramatic folding of the inner nuclear membrane suggesting de novo synthesis of membrane constituents, and concomitantly, accumulation of budding capsids at the inner nuclear membrane as well as fully enveloped virions within the perinuclear space. Shift of virions from the perinuclear space into RER cisternae and lack of foldings of the inner nuclear membrane after Brefeldin A (BFA) treatment suggest that US3 is involved in membrane synthesis needed for envelopment and, concomitantly, in mediating budding and intraluminal virus transportation.

3. Material and Methodes

3.1. Cells and Viruses

Vero cells (European Collection of Cell Cultures) were grown in Dulbecco's modified minimal essential medium (DMEM; Invitrogen, Basel, Switzerland) supplemented with penicillin (100 U/ml), streptomycin (100 µg/ml) and 10% fetal calf serum (FCS; Omnilab, Mettmenstetten, Switzerland). HSV-1 ΔUS3 was kindly provided by Bernard Roizman, University of Chicago, USA. Wild-type HSV-1 strain F and HSV-1 ΔUS3 were propagated in Vero cells. Virus yields were determined by plaque titration.

3.2. Infection of Cells

Vero cells were grown for 2 days on 30 µm thick sapphire disks with a diameter of 3 mm (Bruegger, Minusio, Switzerland) for low temperature electron microscopy (LTEM), or on cover slips for immunolabeling of nuclear pore proteins. Sapphire disks were covered with 8-10 nm carbon, obtained by evaporation under high vacuum conditions, to enhance cell growth. Cells were infected with HSV-1 ΔUS3 at a multiplicity of infection (MOI) of 5 in 1 ml medium per well in 6 well plates, and kept at 37° C for 1 h to allow adsorption prior to incubation at 37°C for 9 to 21 h.

3.3. Brefeldin A

BFA was added in concentrations of 0.1 µl/ml 5 and 8 h after infection of Vero cells with HSV-1 ΔUS3. The total incubation time after inoculation was 20 h.

3.4. Low-Temperature Transmission Electron Microscopy

Cells grown on sapphire disks were frozen in a high-pressure freezing unit (HPM010; BAL-TEC Inc., Balzers, Liechtenstein) as described in detail (Wild, Schraner et al. 2002). The samples were then transferred to a freeze-substitution unit (FS 7500; Boeckeler Instruments, Tucson, Arizona) precooled to -88°C for substitution with acetone and subsequent fixation with 0.25% glutaraldehyde and 0.5% osmium tetroxide at temperatures between -30°C and +2°C to achieve good contrast of membranes (Wild, Schraner et al. 2001), and embedded in Epon at 4°C followed by polymerization at 60°C for 2.5 days. Serial sections of 60 to 90 nm thickness were analyzed in a transmission electron microscope (CM12; Philips, Eindhoven, The Netherlands) equipped with a slow-scan CCD camera (Gatan, Pleasanton, CA) at an acceleration voltage of 100 kV.

3.5. Morphometric Analysis

Images of Golgi complexes were collected from cells 9, 17 and 21 h after infection with HSV-1 Δ US3 at MOI 5 or mock infected cells. Volume density and surface density of the Golgi complex were estimated by morphometric analysis on 30 cells selected at random at a final magnification of 89'500 applying the point counting method (Weibel 1979). The data were calculated on the basis of the equations: $V_{vg} = P_g/P_{cy}$, and $S_{vg} = I_g/(P_{cy} \times d)$, whereby V_{vg} is the volume density, S_{vg} the surface density of the Golgi complex, I_g are intersections with Golgi membranes, P_g are points hitting the lumen of the Golgi complex, P_{cy} are points hitting the cytoplasm, and d is the test line length. The data were expressed as μm^3 volume or as μm^2 surface area per 1000 μm^3 cytoplasm.

3.6. Immunolabeling

For demonstration of nuclear pores by confocal microscopy, cells grown on cover slips were infected with HSV-1 Δ US3 at a MOI 5 and incubated at 37°C. Cells were fixed with 2% formaldehyde in phosphate buffered saline (PBS), pH 7.2, at room temperature (RT) for 25 min, washed in PBS at 4°C for 5 min, incubated with 50 mM Glycine in PBS at 4°C for 5 min, washed in PBS, treated with 0.1% Triton X-100 (E. Merck, Darmstadt) for 5 to 10 min and, after blocking with 3% bovine serum albumin (BSA) in PBST (PBS and 0.05% Tween 20), incubated with MAb 414 (Convance, Berkeley, California, USA) a mouse monoclonal antibody at RT for 1 h. After washing with PBST containing 3% milk, cells were incubated with an anti mouse secondary antibody Alexa Fluor 488 (A 11017; Molecular Probes, Eugene, Oregon, USA). For monitoring virus multiplication cells were then incubated with antibodies against VP16 raised in rabbits (gift from Bernard Roizman, University of Chicago, USA) and finally with a secondary antirabbit antibody conjugated with Alexa Fluor 594 (A 11072, Molecular Probes, Eugene, Oregon, USA). Nuclei were stained with DAPI (Roche). After washing with PBS containing 3% milk and finally with PBS cells were again fixed with 2% formaldehyde for 10 min, washed in water and embedded in Glycergel (DakoCytomation, Carpinteria, USA) with 1,4-Diazabicyclo[2.2.2.] octane (Fluka Chemie GmbH, Buchs, Switzerland). For controls, mock infected cells were used, secondary antibodies were omitted. Samples were analyzed using a confocal laser scanning microscope (SP2, Leica, Mannheim, Germany). Images were deconvolved employing a blind deconvolution algorithm using the program suite Huygens Essential (SVI, Hilversum, The Netherlands).

4. Results

4.1. Nucleus

4.1.1. Nuclear Membranes

Budding of capsids at the inner nuclear membrane is assumed to be inhibited in cells infected with HSV-1 Δ US3 (Wagenaar, Pol et al. 1995; Klupp, Granzow et al. 2000; Reynolds, Ryckman et al. 2001). To clarify the capsid membrane interactions we investigated the nuclear periphery by electron microscopy at improved temporal and spatial resolution. The most impressive alterations found at the nuclear surface were the numerous capsids budding at the inner nuclear membrane, accumulations of virions within the perinuclear space, and the intense folding of the inner nuclear membrane (Fig. 1-3) forming coil-like structures composed of 1 to 6 layers. These coils developed within the nucleus (Fig. 1 and 2A) or protruded into the cytoplasm (Fig. 1 and 2B). Folds within the nucleus consisted exclusively of the inner membranes (Fig. 2A, 3A and B). The outer membrane seems not to be involved in folding. Folds protruding into the cytoplasm consisted probably also of the inner nuclear membrane. They were delineated to the cytoplasm by the outer nuclear membrane (Fig. 2B). The luminal layer of these folded membranes was always thickened by deposition of an electron dense substance. The space between two luminal layers was often filled by a lesser condensed substance than that on the luminal layer.

Capsids were found to bud at the inner nuclear membrane only at sites that were not thickened (Fig. 2A and 3A). The envelope of all virus particles within the perinuclear space contained a similar dense substance (Fig. 3D, 4A and B) as the inner nuclear membrane. Though many of this particles are detached from the nuclear membrane in a given section plane it cannot be ruled out whether they

are still connected to it above or underneath the section plane. The number of virus particles at budding stages and within the perinuclear space was by a factor 2 lower at 20 and 21 hour compared to 17 h suggesting that virions were possibly transported away from the perinuclear space, and consequently, that budding has been completed.




FIG. 1. Tangential section through a nucleus of a Vero cell 19 h after infection with HSV-1 Δ US3 with multiple foldings of nuclear membranes within the nucleus or cytoplasm combined with accumulations of virus particles at various stages of envelopment. Accumulation of capsids (arrows), and dense clusters consisting probably of proteins (p). Bar 1 μ m.

FIG. 2. Vero cells infected with HSV-1 Δ US3 at 17h (A) or 19h (B) post incubation. Bars, 100 nm.

(A) Budding capsid (arrow) at the inner nuclear membrane and folding of the inner nuclear membrane within the nucleus with deposition of a dense substance of unknown nature at the luminal layer (ll) whereas the cytoplasmic layer (cl) is clearly visible.

(B) Foldings of the inner and outer (o) nuclear membrane within the cytoplasm. Virions with a dense envelope are accumulated within the perinuclear space clearly delineated by the inner nuclear membrane (i). The border (arrowheads) between nucleus and cytoplasm is poorly demarcated because the nucleus is tangentially sectioned.





FIG. 3. The nuclear surface of Vero cells 17 or 19 h after infection with HSV-1 Δ US3.
Bars, 100 nm

(A, B) Foldings of the inner nuclear membrane with depositions of an electron dense substance at the luminal layer and budding (arrows) of capsids at various stages at the outermost folding.

(C, D) Accumulation of virions with a dense envelope within the perinuclear space. Capsids bud at the inner nuclear membrane (i). The inner dense nuclear membrane is folded into the cytoplasm together with the outer nuclear membrane (o). The perinuclear space is delineated to the nucleus by the inner nuclear membrane. np: nuclear pore









FIG. 4. Vero cells incubated with HSV-1 Δ US3 at 17 or 19 h. Bars, 100 nm.

(A) Accumulation of virions (v) probably within the perinuclear space, of capsids (c) and probably of proteins (p) within the nuclear matrix.

(B) Accumulations of virions with a dense envelope probably within the perinuclear space as indicated by budding of capsids (arrows). c: capsids;
o: outer nuclear membrane



4.1.2. Nuclear Capsids

Capsids were irregularly scattered within the nucleus often aggregating to clusters. To get an idea about the frequency of capsid clusters we determined the number of clusters per 100 μm^2 by counting them on 25 randomly chosen cellular profiles. In the nuclear matrix about 40 to 50 capsids were found to accumulate to clusters (Fig. 5) often in a quite geometrical manner (Fig. 5D).





FIG. 5. Accumulations of capsids within the nucleus in Vero cells at 17 or 19 h of infection with HSV-1 ΔUS3 . A, B: Bars, 1 μm ; C, D: Bars, 100 nm.

(A) Large clusters of A-, B- and C-capsids within the nuclear matrix and enveloped virions within the perinuclear space clearly delineated by the inner nuclear membrane (i). Foldings of the nuclear membrane (fn) into the cytoplasm, and foldings of the RER (fr).

(B) Mainly B- capsids are scattered throughout the nuclear matrix. Note the undulated dense nuclear envelope.

(C) Accumulation of probably not fully assembled capsid structures surrounded by B-capsids.

(D) Accumulation of mainly B-capsids together with a few A- and C-capsids, and probably unassembled capsid structures.





4.1.3. Nuclear Pores

Most of the nuclear pores detected by electron microscopy were intact. Some nuclear pores, however, were destroyed and nuclear material protruded into cytoplasm but did not merge with it (Fig. 7A). Capsids were not found in association with nuclear pores. Confocal microscopy using antibodies against nuclear pore proteins revealed: that its distribution is irregular in cells infected with HSV-1 Δ US3 compared to mock infected cells (Fig. 6). Infection of cells was recognized by immunolabeling using antibodies against VP 16.

Fig. 6. Confocal imaging of nuclear pores in Vero cells using monoclonal antibodies MAb414. Bar, 3 μ m

(A) Nucleus of mock infected cell after deconvolution showing regular distribution of nuclear pores (green).

(B) Deconvolved nucleus of a Vero cell incubated with HSV-1 Δ US3 for 12 h contains irregularly distributed nuclear pores. Tegument protein VP 16 (red) is expressed in the nucleus and in the cytoplasm.



4.2. Cytoplasm

4.2.1. RER-Golgi complex

Very few virions were found in RER cisternae and in Golgi cisternae (Fig. 7B). The Golgi fields were small at 9 h of incubation (Fig. 7A) and became drastically enlarged at 17 and 21h (Fig. 7B and C) of incubation (Table I). At 17 h p.i. the volume of the Golgi complex was smaller than at 9 and 21 h whereas surface area was higher at 9 and 21 h.

In some Golgi fields we found deposition of proteins at the cytoplasmic layer probably of viral origin (Fig. 7C and D) looking similar to the substance of the viral envelope in the perinuclear space and at the inner nuclear membrane.

Table I: Volume and surface area of the Golgi complex calculated per 1000 μm^3 cytoplasm in cells infected with HSV-1 ΔUS3 at MOI 5

	9 h	17 h	21 h
Surface area	2680	7286	8866
Volume	152	126	153
Surface/ Volume	17.6	57.8	58




FIG. 7. Golgi complex in Vero cells incubated after inoculation with HSV-1 Δ US3 for 9 h, 17 h and 21 h. Bars, 100 nm

(A) Small Golgi field (9 h p.i.) with moderately dilated cisternae adjacent to the nuclear matrix that protruded into the cytoplasm via a markedly dilated (4 μ m) nuclear pore (np) distinctly delineated by the nuclear membranes (i,o).

(B) Large Golgi field (17 h p.i.) that is not involved in envelopment.

(C) Large Golgi field (21 h p.i.) with some deposition of proteins (arrows) most probably of viral origin.

(D) Dispersed Golgi cisternae at 17 h p.i. with deposition of proteins (arrow) probably of viral origin.

4.2.2. Distribution of Phenotypes

In an attempt to get an idea about the distribution of the various phenotypes viral particles within the cytoplasm were counted on images taken at random between 17 and 21 h post inoculation. Table II shows that the majority of phenotypes were at the inner nuclear membrane and within the perinuclear space. The number was by a factor of 2 higher at 17 h compared to 20 and 21 h. The number of naked capsids within the cytoplasm, however, was more than 6 and 5 folds higher, respectively at 20 and 21 h compared to 17 h. The number of virions within vacuoles (Fig. 8) assumed to derive by packaging (P-vacuoles) was low whereas the number of virions within vacuoles assumed to derive by wrapping was about 7 fold higher at 17 and 21 h and even 16 fold higher at 20 h compared to P-vacuoles.

FIG. 8. Virus particles in Vero cells at 19 h of incubation with HSV-1 Δ US3. Bars, 100 nm.

(A) Naked C- and B-capsids within the cytoplasm. Some are involved in budding at membranes of possible Golgi origin. Virions within vacuoles probably derived by wrapping (w) and by packaging (p) for transportation to the plasma membrane.

(B) Large Golgi complex and a laterally dilated cisterna containing one virion with a dense substance at the envelope suggesting its nuclear origin.

p: packaging vacuole

(C) Golgi membranes with a capsid budding at its membrane (b), and a virion within a vacuole probably derived by packaging (p).






Table II: Number of virus particles expressed per 100 μm^2 of cytoplasm in Vero cells incubated for 17, 20 and 21 h after infection with HSV-1 at MOI 5

US3				wt
Phenotypes	17 h	20 h	21 h	17 h
Virus particles at INM and PNS	112.1	60.6	55.1	4
Virions in P vacuoles	2.1	1.8	1.1	4
Nuclear envelopment	114.2	62.4	56.2	8
Naked capsids	4.6	31.2	24.1	3
Virions in W vacuoles	14.9	31.4	14.3	6
Cytoplasmic envelopment	19.5	62.5	38.4	9
Ratio cytoplasmic to nuclear envelopment	1: 5.9	1: 1	1: 1.5	1: 0.9
Nuclear clusters of capsids	4.2	7.4	6.8	1

INM: inner nuclear membrane; **PNS:** perinuclear space; **P-vacuoles:** large vacuoles derived by packaging (Fig. 8); **W-vacuoles:** derived by wrapping: small vacuoles with one virion and an electron-dense substance between envelope and vacuolar membrane.

4.3. Virus Yields

PFU/ml 3×10^7 of infectious virus was determined by plaque titration. The virus yield was 1×10^7 to 3×10^7 similar as in Vero cells infected with wt virus.

4.4. Effects of Brefeldin A on HSV-1 Δ US3 Envelopment

BFA causes disassembly of the Golgi apparatus that rapidly disappears as a morphologically distinct entity (Lippincott-Schwartz, Yuan et al. 1989; Lippincott-Schwartz, Donaldson et al. 1990). Treatment of wt HSV-1 infected cells with BFA was shown to inhibit transportation of virions out of the perinuclear space (Reinhardt 2006). To test whether impaired functionality of the Golgi complex influences envelopment at the inner nuclear membrane we used BFA in concentrations of 1 μ l/ml added at 5 or 8 h post inoculation. Surprisingly, after BFA treatment the budding of capsids at the inner nuclear membrane and the number of virions in the perinuclear space were lower than in untreated cells (Fig. 9) whereas the number of virions within RER cisternae was substantially increased indicating that the virions were transported into RER cisternae (Fig. 10). The foldings of the inner nuclear membrane were less and the outer nuclear membrane was not folded. There was an intense proliferation of RER membranes. The envelope of virions within RER cisternae had a similar dense substance as those in the perinuclear space (Fig. 11).

Fig. 9. Tangential section through a cell nucleus infected with HSV-1 Δ US3 and after treatment with BFA at 8 h p.i. Small foldings of the nuclear membrane within the nucleus (fn) and proliferations of RER membrane (fr). The contour of the nucleus is poorly visible. Most virions (v) got stuck in the perinuclear space-RER compartment. Bar, 1 μ m




Fig 10. Vero cells at 17 h of incubation with HSV-1 Δ US3 and after addition of BFA at 5 h p.i. Bars, 1 μ m.

(A, B) Proliferation of RER membranes (fr) and accumulations of large numbers of virions (v) predominantly within dilated RER cisternae.








Fig. 11. Vero cells infected with HSV-1 Δ US3 for 18 h and after treatment with BFA at 8 h p.i. Bars, 100nm.

(A) Foldings of the RER and dilation of RER cisternae containing enveloped virions.

(B) Accumulation of virions between proliferated RER membranes distinctly thickened by a dense substance.

(C) Proliferation of thickened membranes clearly showing its RER origin.



5. Discussion

High resolution microscopy of HSV-1 Δ US3 infected cells revealed: (i) budding of numerous capsids at the inner nuclear membrane, (ii) accumulations of virions within the perinuclear space, (iii) intense folding of the inner nuclear membrane accompanied by deposition of a dense substance at the luminal layer, (iv) clusters of capsids within the nucleus, (v) dilation of nuclear pores, (vi) enlargement of the Golgi complex with deposition of a dense substance at the cytoplasmic layer, (vii) the presence of naked capsids within the cytoplasm, and (viii) the dominant presence of vacuoles containing a single virion assumed to have originated by budding at Golgi membranes. BFA treatment resulted in (ix) reduced number of budding capsids at the inner nuclear membrane and lower accumulation of virions in the perinuclear space but in (x) accumulation of virions within RER cisternae and (xi) intense proliferation of RER membranes.

A crucial phenomenon in cells infected with HSV-1 Δ US3 is the high number of budding capsids at the inner nuclear membrane and the accumulation of virions in the perinuclear space. Budding of capsids at the inner nuclear membrane is generally accepted as the pathway of capsids to exit the nucleus. The resulting virus particle is a fully enveloped virion containing tegument (Granzow, Klupp et al. 2004; Leuzinger, Ziegler et al. 2005). The high number of budding capsids in HSV-1 Δ US3 infected cells is either caused by retardation of the budding process or by an increased production of capsids. The first idea seems to be more attractive because the onset of budding is much later in HSV-1 Δ US3 compared to wt HSV-1 infected cells that occurred as early as 6 h p.i. (Leuzinger, Ziegler et al. 2005) and because capsids also accumulated within the nucleus often to large clusters. Accumulation of virions within the perinuclear space of cells infected with HSV-1 Δ US3 was already described (Reynolds, Wills et al. 2002) whereas very recently these accumulations has been suggested

to be within vacuoles formed by the inner nuclear membrane (Poon, Benetti et al. 2006). We clearly show that the inner nuclear membrane extended deeply into the nucleus (Fig. 1) giving the impression of vacuoles (Fig. 3) depending on the section plane. The outer nuclear membrane, however, is not involved in the formation of these foldings.

Accumulation of virions within the perinuclear space indicates that transportation out of it is inhibited. In the absence of the PrV US3 protein homologue, large numbers of enveloped virions appeared to accumulate within invaginations of the nuclear membrane. This observation led to the deduction that the US3 encoded kinase is important for the efficient de-envelopment of nascent virions by fusion of the viral envelope with the outer nuclear membrane (Wagenaar, Pol et al. 1995; Klupp 2001). We found a low number of virions within Golgi cisternae indicating that a few virions were intraluminally transported out of the perinuclear space. We, therefore, exposed cells to BFA added 5 or 8 h post inoculation with HSV-1 Δ US3. Surprisingly, treatment with BFA resulted in a reduction of budding capsids at the inner nuclear membrane and of virions within the perinuclear space whereas the number of virions in adjacent RER cisternae was substantially increased clearly indicating that virions exit the perinuclear space by intraluminal transportation as have been shown for wildtype HSV-1 (Schwartz and Roizman 1969; Whealy, Card et al. 1991; Gilbert, Ghosh et al. 1994; Radsak, Eickmann et al. 1996; Granzow, Weiland et al. 1997; Roller, Zhou et al. 2000; Leuzinger, Ziegler et al. 2005). The Golgi complex disintegrates within minutes after administration of BFA to cell cultures (Hess, Muller et al. 2000). The regained ability of virions due to the lack of US3 for intraluminal transportation after dissociation of Golgi membranes suggests that the primary action site of US3 kinase is at the level of the Golgi complex. This idea is strongly supported by the fact that the foldings at the inner nuclear membrane is drastically reduced in HSV-1 Δ US3 infected

cells after treatment with BFA indicating that US3 is involved in the regulation of distribution and possibly of synthesis of membrane constituents required for envelopment, possibly in concert with facilitation of viral transport out of the perinuclear space. De novo synthesized phospholipids have been shown to be inserted into nuclear membranes of PrV infected cells (Ben-Porat and Kaplan 1971; Ben-Porat and Kaplan 1972) and HSV-1 infected cells (Sutter 2006). It is thus reasonable to assume that the dramatic foldings of the inner nuclear membrane is the result of enhanced phospholipid synthesis and/or retarded budding resulting in an imbalance of supply and demand of membrane constituent for envelopment. This seems to be also true for membranes of the Golgi complex which becomes enlarged by a factor of about 3 at 17 and 21 h respectively, compared to 9 h of infection. In the context of the de-envelopment theory this substantial enlargement would be explained by the lack of need of membranes for wrapping because all the virions got stuck in the perinuclear space. Our data clearly show that capsids got access to the cytoplasm. 1/3 to 3/4 of capsids—depending on the incubation time—have to acquire tegument and an envelope by budding at Golgi membranes. If the lack of US3 inhibits de-envelopment (Klupp 2001; Reynolds, Wills et al. 2002; Granzow, Klupp et al. 2004) these naked capsids within the cytoplasm will have to take another route from the nucleus to the cytoplasm then via fusion at the outer nuclear membrane. The only alternative on the basis of the current knowledge is via impaired nuclear pores as reported for HSV-1 (Leuzinger, Ziegler et al. 2005) and BHV-1 (Wild, Schraner et al. 2002).

Budding capsids at the inner nuclear membrane was shown to start by thickening of the membrane (Wild, Schraner et al. 2002; Leuzinger, Ziegler et al. 2005) due to deposition of an electron dense substance. This substance remains on the viral envelope within the perinuclear space. The significance of this substance is speculated to be responsible for the ability of intraluminal

transportation (Leuzinger, Ziegler et al. 2005), possibly, in combination with UL31 and UL 34. UL31 and UL34 proteins form a complex that accumulates at the nuclear membrane. It is assumed that this protein complex plays an important role in envelopment of capsids (Reynolds, Ryckman et al. 2001). US3 is suggested to be required for the distribution of UL31 and UL34 proteins throughout the nuclear rim (Reynolds, Ryckman et al. 2001). One possibility is that UL34 protein interacts directly with capsids and/or tegument components at the nuclear membranes (Reynolds, Ryckman et al. 2001). The UL34 protein is possibly responsible for recruiting other viral or cellular factors to the site of envelopment. Other proteins including gK and UL11 are implicated in the initial budding of herpesvirus capsids at the inner nuclear membrane (Baines and Roizman 1992). It is thus reasonable to assume that the dense substance at the viral envelope in the perinuclear space is a complex of UL11, UL31, UL34, gK and possibly other substances.

The question arises whether the substances incorporated into the folded inner nuclear membrane in HSV-1 Δ US3 infected cells are identical to that of the viral envelope in the perinuclear space. Interestingly, the dense substance at the nuclear foldings seems always to be inserted at the luminal layer. Budding takes place at the cytoplasmic site that becomes thickened during the budding process. A very early event in budding is thickening of the membrane at the budding front by insertion of an electron dense substance starting at the luminal layer suggesting that the mechanism of deposition is similar and possibly the composition as well. To clarify the nature of the substances involved in budding on the one hand, and in thickening of the folded inner nuclear membrane on the other, immunolabeling studies at the electron microscopic level must be performed.

Large amounts of dense substances were found also to be deposited at the cytoplasmic layer of the Golgi complex filling the space between two adjacent cytoplasmic layers, i.e. exactly at the opposite layer as at the inner nuclear membrane. The difference in location to the substances at the inner nuclear membrane and at the viral envelope in the perinuclear space suggests different composition and function of the substance. Possibly, the bulk of the substance consists of tegument proteins that need also to be investigated by immunogoldlabeling.

After treatment with BFA there were intense proliferation of RER membranes whereas nuclear foldings were almost absent. A dense substance was deposited between two adjacent cytoplasmic layers similar as between Golgi membranes. Dissociated Golgi membranes are rearranged in the RER compartment (Lippincott-Schwartz, Yuan et al. 1989; Lippincott-Schwartz, Donaldson et al. 1990). We, therefore, assume that the proliferation of RER membranes is due to a membrane shift from the Golgi complex, possibly together with the dense substances. Immunolabeling studies will probably shed some light to the origin of these substances.

US3 is not essential for the production of infectious virions (Reynolds, Wills et al. 2002). Compared to wt infected cells, the virus yield was similar in the range of $1-3 \times 10^7$ PFU/ml. The most interesting question is thus where do all these virions that cannot escape the perinuclear space (Reynolds, Wills et al. 2002) or nuclear vacuoles (Poon, Benetti et al. 2006) acquire the requirements for infectivity? The simple answer is: in the course of budding at the inner nuclear membrane. According to table I 76 to 88% of all enveloped virus particles were within the perinuclear space. Consequently, the majority of virions contributing to the yield of infectious virus had originated by budding at the inner nuclear membrane. That means that virions acquire all essential components including

tegument proteins and glycoproteins by budding at the inner nuclear membrane as suggested (Stannard, Himmelhoch et al. 1996). Indeed, many glycoproteins have been shown to be present at nuclear membranes (Hutchinson, Roop-Beauchamp et al. 1995; Rychlowski, Rijsewijk et al. 2001). Consequently, budding of capsids leads to an enveloped virion containing glycoproteins. The deposition of the dense substance of the viral envelope impressively demonstrable in virions within the perinuclear space must, hence, been considered to cover the glycoproteins so that they cannot be detected by immunolabeling (Stannard, Himmelhoch et al. 1996). The significance of this dense substance was considered to facilitate transportation of virions out of the perinuclear space (Pol, Wagenaar et al. 1991; Leuzinger, Ziegler et al. 2005). One of these substances contributing directly or indirectly can be assumed to be US3.

In conclusion, the data obtained by high resolution microscopy confirmed the idea of retarded envelopment at the inner nuclear membrane in US3 deleted HSV-1 mutants. Furthermore, they clearly show, i) that large amounts of membranes are needed for envelope formation by budding at the inner nuclear membrane which is not inserted into the outer nuclear membrane, ii) that virions are transported from the perinuclear space into RER cisternae, iii) that a minor proportion of capsids gain access to the cytoplasm probably via dilated pores, iv) that virions get stuck in the perinuclear space after budding of capsids at the inner nuclear membrane is completed, and v) budding of capsids at Golgi membranes is retarded. It, hence, can be concluded that the bulk of virions contributing to the total infectious virus yield originates at the inner nuclear membrane, and furthermore, that US3 is involved in the maintenance of envelope formation, possibly in concert with intraluminal virus transportation.

6. References

- Asano, S., T. Honda, et al. (1999). "US3 protein kinase of herpes simplex virus type 2 plays a role in protecting corneal epithelial cells from apoptosis in infected mice." Journal of General Virology **80**(Pt 1): 51-6.
- Avitabile, E., P. L. Ward, et al. (1994). "The herpes simplex virus UL20 protein compensates for the differential disruption of exocytosis of virions and viral membrane glycoproteins associated with fragmentation of the Golgi apparatus." Journal of Virology **68**(11): 7397-405.
- Baer, R., A. T. Bankier, et al. (1984). "DNA sequence and expression of the B95-8 Epstein-Barr virus genome." Nature **310**(5974): 207-11.
- Baines, J. D. and B. Roizman (1992). "The UL11 gene of herpes simplex virus 1 encodes a function that facilitates nucleocapsid envelopment and egress from cells." Journal of Virology **66**(8): 5168-74.
- Baines, J. D., P. L. Ward, et al. (1991). "The UL20 gene of herpes simplex virus 1 encodes a function necessary for viral egress." Journal of Virology **65**(12): 6414-24.
- Ben-Porat, T. and A. S. Kaplan (1971). "Phospholipid metabolism of herpesvirus-infected and uninfected rabbit kidney cells." Virology **45**(1): 252-64.
- Ben-Porat, T. and A. S. Kaplan (1972). "Studies on the biogenesis of herpesvirus envelope." Nature **235**(5334): 165-6.
- Blaho, J. A., C. Mitchell, et al. (1994). "An amino acid sequence shared by the herpes simplex virus 1 alpha regulatory proteins 0, 4, 22, and 27 predicts the nucleotidylation of the UL21, UL31, UL47, and UL49 gene products." Journal of Biological Chemistry **269**(26): 17401-10.
- Campadelli-Fiume, G., F. Farabegoli, et al. (1991). "Origin of unenveloped capsids in the cytoplasm of cells infected with herpes simplex virus 1." Journal of Virology **65**(3): 1589-95.
- Chang, Y. E. and B. Roizman (1993). "The product of the UL31 gene of herpes simplex virus 1 is a nuclear phosphoprotein which partitions with the nuclear matrix." Journal of Virology **67**(11): 6348-56.
- Chang, Y. E., C. Van Sant, et al. (1997). "The null mutant of the U(L)31 gene of herpes simplex virus 1: construction and phenotype in infected cells." Journal of Virology **71**(11): 8307-15.
- Chee, M. S., A. T. Bankier, et al. (1990). "Analysis of the protein-coding content of the sequence of human cytomegalovirus strain AD169." Current Topics in Microbiology & Immunology **154**: 125-69.
- Darlington, R. W. and L. H. d. Moss (1968). "Herpesvirus envelopment." Journal of Virology **2**(1): 48-55.
- Di Lazzaro, C., G. Campadelli-Fiume, et al. (1995). "Intermediate forms of glycoconjugates are present in the envelope of herpes simplex virions during their transport along the exocytic pathway." Virology **214**(2): 619-23.
- Foster, T. P. and K. G. Kousoulas (1999). "Genetic analysis of the role of herpes simplex virus type 1 glycoprotein K in infectious virus production and egress." Journal of Virology **73**(10): 8457-68.
- Foster, T. P., G. V. Rybachuk, et al. (2003). "Overexpression of gK in gK-transformed cells collapses the Golgi apparatus into the endoplasmic reticulum inhibiting virion egress, glycoprotein transport, and virus-induced cell fusion." Virology **317**(2): 237-52.
- Frame, M. C., F. C. Purves, et al. (1987). "Identification of the herpes simplex virus protein kinase as the product of viral gene US3." Journal of General Virology **68**(Pt 10): 2699-704.

- Fuchs, W., B. G. Klupp, et al. (1997). "The UL20 gene product of pseudorabies virus functions in virus egress." Journal of Virology **71**(7): 5639-46.
- Fuchs, W., B. G. Klupp, et al. (2002). "The interacting UL31 and UL34 gene products of pseudorabies virus are involved in egress from the host-cell nucleus and represent components of primary enveloped but not mature virions." Journal of Virology **76**(1): 364-78.
- Galvan, V. and B. Roizman (1998). "Herpes simplex virus 1 induces and blocks apoptosis at multiple steps during infection and protects cells from exogenous inducers in a cell-type-dependent manner." Proceedings of the National Academy of Sciences of the United States of America **95**(7): 3931-6.
- Gershon, A. A., D. L. Sherman, et al. (1994). "Intracellular transport of newly synthesized varicella-zoster virus: final envelopment in the trans-Golgi network." Journal of Virology **68**(10): 6372-90.
- Gibson, W. (1996). "Structure and assembly of the virion." Intervirology **39**(5-6): 389-400.
- Gilbert, R. and H. P. Ghosh (1993). "Immunoelectron microscopic localization of herpes simplex virus glycoprotein gB in the nuclear envelope of infected cells." Virus Research **28**(3): 217-31.
- Gilbert, R., K. Ghosh, et al. (1994). "Membrane anchoring domain of herpes simplex virus glycoprotein gB is sufficient for nuclear envelope localization." Journal of Virology **68**(4): 2272-85.
- Granzow, H. (2001). "Egress of alphaherpesviruses: comparative ultrastructural study." Journal of Virology **75**(8): 3675-84.
- Granzow, H., B. G. Klupp, et al. (2004). "The pseudorabies virus US3 protein is a component of primary and of mature virions." Journal of Virology **78**(3): 1314-23.
- Granzow, H., F. Weiland, et al. (1997). "Ultrastructural analysis of the replication cycle of pseudorabies virus in cell culture: a reassessment." Journal of Virology **71**(3): 2072-82.
- Harper, D. R. and H. O. Kangro (1990). "Lipoproteins of varicella-zoster virus." Journal of General Virology **71**(Pt 2): 459-63.
- Hess, M. W., M. Muller, et al. (2000). "Cryopreparation provides new insight into the effects of brefeldin A on the structure of the HepG2 Golgi apparatus." Journal of Structural Biology **130**(1): 63-72.
- Hutchinson, L., C. Roop-Beauchamp, et al. (1995). "Herpes simplex virus glycoprotein K is known to influence fusion of infected cells, yet is not on the cell surface." Journal of Virology **69**(7): 4556-63.
- Jerome, K. R., R. Fox, et al. (1999). "Herpes simplex virus inhibits apoptosis through the action of two genes, Us5 and Us3." Journal of Virology **73**(11): 8950-7.
- Jones, F. and C. Grose (1988). "Role of cytoplasmic vacuoles in varicella-zoster virus glycoprotein trafficking and virion envelopment." Journal of Virology **62**(8): 2701-11.
- Klupp, B. G. (2001). "Effect of the pseudorabies virus US3 protein on nuclear membrane localization of the UL34 protein and virus egress from the nucleus." Journal of General Virology **82**(Pt 10): 2363-71.
- Klupp, B. G. (2001). "Pseudorabies virus UL37 gene product is involved in secondary envelopment." Journal of Virology **75**(19): 8927-36.
- Klupp, B. G., W. Fuchs, et al. (2002). "Pseudorabies virus UL36 tegument protein physically interacts with the UL37 protein." Journal of Virology **76**(6): 3065-71.
- Klupp, B. G., H. Granzow, et al. (2000). "Primary envelopment of pseudorabies virus at the nuclear membrane requires the UL34 gene product." Journal of Virology **74**(21): 10063-10073.
- Klupp, B. G., R. Nixdorf, et al. (2000). "Pseudorabies virus glycoprotein M inhibits membrane fusion." Journal of Virology **74**(15): 6760-8.

- Kopp, M., H. Granzow, et al. (2004). "Simultaneous deletion of pseudorabies virus tegument protein UL11 and glycoprotein M severely impairs secondary envelopment." Journal of Virology **78**(6): 3024-34.
- Kopp, M., H. Granzow, et al. (2003). "The pseudorabies virus UL11 protein is a virion component involved in secondary envelopment in the cytoplasm." Journal of Virology **77**(9): 5339-51.
- Kopp, M., B. G. Klupp, et al. (2002). "Identification and characterization of the pseudorabies virus tegument proteins UL46 and UL47: role for UL47 in virion morphogenesis in the cytoplasm." Journal of Virology **76**(17): 8820-33.
- Lee, K., J. Bao, et al. (1987). "Ultrastructural study of the morphogenesis of herpes simplex virus type 2 in organ cultured human uterine cervix and the interaction between virus and host cell." Journal of Electron Microscopy Technique **7**: 73-84.
- Leopardi, R., C. Van Sant, et al. (1997). "The herpes simplex virus 1 protein kinase US3 is required for protection from apoptosis induced by the virus." Proceedings of the National Academy of Sciences of the United States of America **94**(15): 7891-6.
- Leuzinger, H., U. Ziegler, et al. (2005). "Herpes simplex virus 1 envelopment follows two diverse pathways." Journal of Virology **79**: 13047-13059.
- Lippincott-Schwartz, J., J. G. Donaldson, et al. (1990). "Microtubule-dependent retrograde transport of proteins into the ER in the presence of brefeldin A suggests an ER recycling pathway." Cell **60**(5): 821-36.
- Lippincott-Schwartz, J., L. C. Yuan, et al. (1989). "Rapid redistribution of Golgi proteins into the ER in cells treated with brefeldin A: evidence for membrane cycling from Golgi to ER." Cell **56**(5): 801-13.
- Loomis, J. S., J. B. Bowzard, et al. (2001). "Intracellular trafficking of the UL11 tegument protein of herpes simplex virus type 1." Journal of Virology **75**(24): 12209-19.
- MacLean, C. A., B. Clark, et al. (1989). "Gene UL11 of herpes simplex virus type 1 encodes a virion protein which is myristylated." Journal of General Virology **70**(Pt 12): 3147-57.
- McMillan, T. N. and D. C. Johnson (2001). "Cytoplasmic domain of herpes simplex virus gE causes accumulation in the trans-Golgi network, a site of virus envelopment and sorting of virions to cell junctions." Journal of Virology **75**(4): 1928-40.
- Mettenleiter, T. C. (2002). "Herpesvirus assembly and egress." Journal of Virology **76**(4): 1537-47.
- Mettenleiter, T. C. (2004). "Budding events in herpesvirus morphogenesis." Virus Research **106**(2): 167-80.
- Morgan, C., S. A. Ellison, et al. (1954). "Structure and development of viruses as observed in the electron microscope." Journal of Experimental Medicine **100**: 195-202.
- Mossman, K. L., R. Sherburne, et al. (2000). "Evidence that herpes simplex virus VP16 is required for viral egress downstream of the initial envelopment event." Journal of Virology **74**(14): 6287-99.
- Munger, J., A. V. Chee, et al. (2001). "The U(S)3 protein kinase blocks apoptosis induced by the d120 mutant of herpes simplex virus 1 at a premitochondrial stage." Journal of Virology **75**(12): 5491-7.
- Muranyi, W., J. Haas, et al. (2002). "Cytomegalovirus recruitment of cellular kinases to dissolve the nuclear lamina.[see comment]." Science **297**(5582): 854-7.
- Pol, J. M., F. Wagenaar, et al. (1991). "Morphogenesis of three pseudorabies virus strains in porcine nasal mucosa." Intervirology **32**(6): 327-37.
- Poliquin, L., G. Levine, et al. (1985). "Involvement of Golgi apparatus and a restructured nuclear envelope during biogenesis and transport of herpes simplex virus glycoproteins." Journal of Histochemistry & Cytochemistry **33**(9): 875-83.

- Poon, A. P., L. Benetti, et al. (2006). "U(S)3 and U(S)3.5 protein kinases of herpes simplex virus 1 differ with respect to their functions in blocking apoptosis and in virion maturation and egress." Journal of Virology **80**(8): 3752-64.
- Purves, F. C., A. D. Deana, et al. (1986). "The substrate specificity of the protein kinase induced in cells infected with herpesviruses: studies with synthetic substrates [corrected] indicate structural requirements distinct from other protein kinases.[erratum appears in Biochim Biophys Acta 1987 Feb 18;927(2):303]." Biochimica et Biophysica Acta **889**(2): 208-15.
- Purves, F. C., M. Katan, et al. (1986). "Characteristics of the induction of a new protein kinase in cells infected with herpesviruses." Journal of General Virology **67**(Pt 6): 1049-57.
- Purves, F. C., R. M. Longnecker, et al. (1987). "Herpes simplex virus 1 protein kinase is encoded by open reading frame US3 which is not essential for virus growth in cell culture." Journal of Virology **61**(9): 2896-901.
- Purves, F. C. and B. Roizman (1992). "The UL13 gene of herpes simplex virus 1 encodes the functions for posttranslational processing associated with phosphorylation of the regulatory protein alpha 22." Proceedings of the National Academy of Sciences of the United States of America **89**(16): 7310-4.
- Purves, F. C., D. Spector, et al. (1991). "The herpes simplex virus 1 protein kinase encoded by the US3 gene mediates posttranslational modification of the phosphoprotein encoded by the UL34 gene." Journal of Virology **65**(11): 5757-64.
- Purves, F. C., D. Spector, et al. (1992). "UL34, the target of the herpes simplex virus U(S)3 protein kinase, is a membrane protein which in its unphosphorylated state associates with novel phosphoproteins." Journal of Virology **66**(7): 4295-303.
- Radsak, K., M. Eickmann, et al. (1996). "Retrieval of human cytomegalovirus glycoprotein B from the infected cell surface for virus envelopment." Archives of Virology **141**(3-4): 557-72.
- Reinhardt, L. (2006). "Significance of the Golgi apparatus in envelopment of herpes simplex virus 1." Thesis, Zurich.
- Reynolds, A. E., B. J. Ryckman, et al. (2001). "U(L)31 and U(L)34 proteins of herpes simplex virus type 1 form a complex that accumulates at the nuclear rim and is required for envelopment of nucleocapsids." Journal of Virology **75**(18): 8803-17.
- Reynolds, A. E., E. G. Wills, et al. (2002). "Ultrastructural localization of the herpes simplex virus type 1 UL31, UL34, and US3 proteins suggests specific roles in primary envelopment and egress of nucleocapsids." Journal of Virology **76**(17): 8939-52.
- Roffman, E., J. P. Albert, et al. (1990). "Putative site for the acquisition of human herpesvirus 6 virion tegument." Journal of Virology **64**(12): 6308-13.
- Roller, R. J., Y. Zhou, et al. (2000). "Herpes simplex virus type 1 U(L)34 gene product is required for viral envelopment." Journal of Virology **74**(1): 117-29.
- Rychlowski, M., F. A. M. Rijsewijk, et al. (2001). "Tyrosine 467 in cytoplasmic tail of glycoprotein E of bovine herpesvirus 1 is essential for gE/gI intracellular trafficking." Proc. of 26th Int. Herpesvirus Workshop, 2001, abstr. 2.27.
- Sanchez, V., E. Sztul, et al. (2000). "Human cytomegalovirus pp28 (UL99) localizes to a cytoplasmic compartment which overlaps the endoplasmic reticulum-golgi-intermediate compartment." Journal of Virology **74**(8): 3842-51.
- Schimmer, C. and A. Neubauer (2003). "The equine herpesvirus 1 UL11 gene product localizes to the trans-golgi network and is involved in cell-to-cell spread." Virology **308**(1): 23-36.
- Schwartz, J. and B. Roizman (1969). "Concerning the egress of herpes simplex virus from infected cells: electron and light microscope observations." Virology **38**(1): 42-9.

- Scott, E. S. and P. O'Hare (2001). "Fate of the inner nuclear membrane protein lamin B receptor and nuclear lamins in herpes simplex virus type 1 infection." Journal of Virology **75**(18): 8818-30.
- Severi, B., M. P. Landini, et al. (1988). "Human cytomegalovirus morphogenesis: an ultrastructural study of the late cytoplasmic phases." Archives of Virology **98**(1-2): 51-64.
- Shiba, C., T. Daikoku, et al. (2000). "The UL34 gene product of herpes simplex virus type 2 is a tail-anchored type II membrane protein that is significant for virus envelopment." Journal of General Virology **81**(Pt 10): 2397-405.
- Silva, M. C., Q. C. Yu, et al. (2003). "Human cytomegalovirus UL99-encoded pp28 is required for the cytoplasmic envelopment of tegument-associated capsids." Journal of Virology **77**(19): 10594-605.
- Siminoff, P. and M. G. Menefee (1966). "Normal and 5-bromodeoxyuridine-inhibited development of herpes simplex virus. An electron microscope study." Experimental Cell Research **44**(2): 241-55.
- Skepper, J. N., A. Whiteley, et al. (2001). "Herpes simplex virus nucleocapsids mature to progeny virions by an envelopment --> deenvelopment --> reenvelopment pathway." Journal of Virology **75**(12): 5697-702.
- Stackpole, C. W. (1969). "Herpes-type virus of the frog renal adenocarcinoma. I. Virus development in tumor transplants maintained at low temperature." Journal of Virology **4**(1): 75-93.
- Stannard, L. M., S. Himmelhoch, et al. (1996). "Intra-nuclear localization of two envelope proteins, gB and gD, of herpes simplex virus." Archives of Virology **141**(3-4): 505-24.
- Steven, A., , Spear, P G (1997). Herpesvirus capsid assembly and envelopment. Structural Biology of viruses. W. Chiu, Burnett, R M, Garcea, R L. New York, Oxford, Oxford University Press: 312-351.
- Sutter, E. (2006). "Phospholipid Metabolism of HSV-1 Infected Cells." Thesis, Zürich.
- Tooze, J., M. Hollinshead, et al. (1993). "Progeny vaccinia and human cytomegalovirus particles utilize early endosomal cisternae for their envelopes." European Journal of Cell Biology **60**(1): 163-78.
- Torrisi, M. R., C. Di Lazzaro, et al. (1992). "Herpes simplex virus envelopment and maturation studied by fracture label." Journal of Virology **66**(1): 554-61.
- van Drunen Littel-van den Hurk, S., S. Garzon, et al. (1995). "The role of the major tegument protein VP8 of bovine herpesvirus-1 in infection and immunity." Virology **206**(1): 413-25.
- van Genderen, I. L., R. Brandimarti, et al. (1994). "The phospholipid composition of extracellular herpes simplex virions differs from that of host cell nuclei." Virology **200**(2): 831-6.
- Wagenaar, F., J. M. Pol, et al. (1995). "The US3-encoded protein kinase from pseudorabies virus affects egress of virions from the nucleus." Journal of General Virology **76**(Pt 7): 1851-9.
- Ward, P. L., G. Campadelli-Fiume, et al. (1994). "Localization and putative function of the UL20 membrane protein in cells infected with herpes simplex virus 1." Journal of Virology **68**(11): 7406-17.
- Weibel, E. (1979). Stereological methods, Vol.I. Practical methods for biological morphometry, Academic Press, London.
- Whealy, M. E., J. P. Card, et al. (1991). "Effect of brefeldin A on alphaherpesvirus membrane protein glycosylation and virus egress." Journal of Virology **65**(3): 1066-81.
- Whiteley, A., B. Bruun, et al. (1999). "Effects of targeting herpes simplex virus type 1 gD to the endoplasmic reticulum and trans-Golgi network." Journal of Virology **73**(11): 9515-20.

- Wild, P., M. Engels, et al. (2005). "Impairment of nuclear pores in bovine herpesvirus 1-infected MDBK cells." Journal of Virology **79**(2): 1071-83.
- Wild, P., E. M. Schraner, et al. (2001). "Enhanced resolution of membranes in cultured cells by cryoimmobilization and freeze-substitution." Microscopy Research and Technique.
- Wild, P., E. M. Schraner, et al. (2002). "The significance of the Golgi complex in envelopment of bovine herpesvirus 1 (BHV-1) as revealed by cryobased electron microscopy." Micron **33**(4): 327-37.
- Yamauchi, Y., C. Shiba, et al. (2001). "Herpes simplex virus type 2 UL34 protein requires UL31 protein for its relocation to the internal nuclear membrane in transfected cells." Journal of General Virology **82**(Pt 6): 1423-8.
- Ye, G. J. and B. Roizman (2000). "The essential protein encoded by the UL31 gene of herpes simplex virus 1 depends for its stability on the presence of UL34 protein." Proceedings of the National Academy of Sciences of the United States of America **97**(20): 11002-7.
- Zhou, Z. H., D. H. Chen, et al. (1999). "Visualization of tegument-capsid interactions and DNA in intact herpes simplex virus type 1 virions." Journal of Virology **73**(4): 3210-8.
- Zhu, Z., M. D. Gershon, et al. (1995). "Envelopment of varicella-zoster virus: targeting of viral glycoproteins to the trans-Golgi network." Journal of Virology **69**(12): 7951-9.

CURRICULUM VITAE

Personalien

Name: Leisinger
Vorname: Sabine
Geburtsdatum: 7. April 1979
Heimatort: Davos
Nationalität: Schweiz
Geburtsort: Dornach

Ausbildung

2006	Doktorandin am Veterinäranatomischen Institut der Vetsuisse Fakultät Zürich
1999-05	Studium der Veterinärmedizin an der Universität Zürich
1995-99	Schweizerische Alpine Mittelschule Davos mit Maturaabschluss Typus E
1992-95	Sekundarschule Davos Platz
1986-92	Primarschule Davos Dorf

ACKNOWLEDGEMENTS

Ich möchte mich ganz herzlich bei allen bedanken, die mich in irgendeiner Form bei meiner Arbeit unterstützt haben. Mein besonderer Dank gilt:

Prof. Dr. P. Wild, für die Überlassung des Dissertationsthemas, für seine sehr gute Unterstützung und den Einblick in den wissenschaftlichen Bereich,

Prof. Dr. M. Ackermann, für die Übernahme des Korreferats,

Elisabeth Schraner, für die Tips und Tricks beim Arbeiten im Labor, ihre Geduld und die professionelle Hilfe bei computertechnischen Problemen,

Elisabeth Högger-Manser, für die Einführung des Ultradünnschneidens,

den **Mitarbeitern des Virologischen Instituts der Vetsuisse Fakultät Zürich**, unter der Leitung von Prof. M. Ackermann. Speziellen Dank an **Claudia Senn** für ihre fachliche Unterstützung,

meinen Mitdoktorandinnen **Céline Manera** und **Ladina Reinhardt**, für die unterhaltsamen Gespräche während und neben der Arbeit,

meinen **Eltern**, für ihre grossartige und liebevolle Unterstützung in den letzten Jahren,

und zuletzt ein Dankeschön an meinen Freund, meine Brüder, meine Schwägerinnen und allen meinen Freunden für ihre tatkräftige Unterstützung.